



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁴ : C12N 15/00, 7/00, A61K 39/285 A61K 39/21, 39/145, 45/02 A61K 37/02 // C12R 1:91	A1	(11) International Publication Number: WO 88/00971 (43) International Publication Date: 11 February 1988 (11.02.88)
--	----	--

(21) International Application Number: PCT/AU87/00246

(22) International Filing Date: 31 July 1987 (31.07.87)

(31) Priority Application Number: PH 07212/86

(32) Priority Date: 1 August 1986 (01.08.86)

(33) Priority Country: AU

(71) Applicants (for all designated States except US): COMMONWEALTH SCIENTIFIC AND INDUSTRIAL RESEARCH ORGANISATION [AU/AU]; Limestone Avenue, Campbell, ACT 2601 (AU). THE AUSTRALIAN NATIONAL UNIVERSITY [AU/AU]; Acton, ACT 2601 (AU).

(72) Inventors; and

(75) Inventors/Applicants (for US only): RAMSHAW, Ian, Allister [AU/AU]; 28 Kallara Close, Duffy, ACT 2611 (AU). BOYLE, David, Bernard [AU/AU]; 3/244 Myers Street, Geelong, VIC 3220 (AU). COUPAR, Barbara, Elizabeth, Howieson [GB/AU]; 12 Thear Street, East Geelong, VIC 3219 (AU). ANDREW, Marion, Elizabeth [AU/AU];

36 Henry Street, Belmont, VIC 3216 (AU).

(74) Agents: SLATTERY, John, Michael et al.; Davies & Collison, 1 Little Collins Street, Melbourne, VIC 3000 (AU).

(81) Designated States: AT (European patent), AU, BE (European patent), CH (European patent), DE (European patent), FR (European patent), GB (European patent), IT (European patent), JP, KR, LU (European patent), NL (European patent), SE (European patent), US.

Published


With international search report.

Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.

(54) Title: RECOMBINANT VACCINE

(57) Abstract

A recombinant vaccine comprises a vaccine vector which incorporates a first nucleotide sequence capable of being expressed as all or a part of an antigenic polypeptide, together with a second nucleotide sequence capable of being expressed as all or a part of a lymphokine effective in enhancing the immune response to the antigenic polypeptide. The vaccine vectors include poxvirus, herpes virus or adenovirus, and the lymphokine may be an interleukin or gamma-interferon. The vaccine vector may express an antigenic polypeptide which is foreign to the host vector.



"RECOMBINANT VACCINE"

This invention concerns new vaccines developed using recombinant DNA technology to provide useful immune responses in circumstances where traditional vaccines may not be sufficiently
5 effective.

Many existing live or killed vaccines are not without disadvantages, often significant, in respect of, for example, high production costs, poor response, low response to poorly immunogenic
10 antigens, instability and a requirement for adjuvants. Furthermore, alternative vaccine preparations based on agents such as purified proteins or synthetic peptide antigens frequently offer only poor protection. In response to these
15 problems, attention has turned to the development of vaccines in which recombinant DNA methods have been used to introduce antigens to which immunity is required, into carrier viruses such as vaccinia.

The advantage of the recombinant DNA approach is that an infectious recombinant virus simultaneously synthesizes the foreign polypeptide and viral antigen, which can then be delivered to a host immune system as a superficial skin lesion. Vaccinia viruses have, for example, been modified for expression of the genes for hepatitis B, human immunodeficiency virus, influenza and malaria antigens; the construction of recombinant viruses carrying other antigens of medical or veterinary importance is under investigation.

In some instances, however, the immune response of recombinant vaccines may be of limited nature and magnitude. Thus, while peripheral immunization with vaccinia-influenza recombinants provides good protection against lower respiratory tract infection, it fails to induce immunity in the upper respiratory tract. On the other hand, peripheral immunization with recombinant vaccines may prove ineffective when local rather than systemic immunity is required, as in say the gastro-intestinal tract.

There have been various attempts to remedy these deficiencies, including expression of vaccine antigens through viruses having stronger promoters, such as poxvirus, but to date these have not met with significant success. The present invention provides an effective means for enhancing the immune response to the specific foreign antigenic polypeptides of recombinant vaccines.

The immune system is regulated in part by molecules, known as lymphokines, which are released by lymphocytes and help or modify the functions of other classes of lymphocytes. The present invention

is based on a recognition that the expression of appropriate lymphokines from recombinant bacterial or viral vaccines can boost and/or modify the immune response to viral, bacterial or co-expressed foreign antigenic polypeptides.

Accordingly, in its broadest aspect, this invention provides a recombinant vaccine comprising a vaccine vector which incorporates a first nucleotide sequence capable of being expressed as all or a part of an antigenic polypeptide, together with a second nucleotide sequence capable of being expressed as all or a part of a lymphokine effective in enhancing the immune response to the antigenic polypeptide.

In accordance with one embodiment of this invention, the first nucleotide sequence capable of being expressed as all or a part of an antigenic polypeptide may be a "native" sequence of the host vector itself, for example vaccinia or herpes virus. In this embodiment, administration of the vaccine of this invention provides augmentation and/or selective induction of the immune response to the "native" antigenic polypeptide.

In another embodiment, the vaccine vector incorporates a nucleotide sequence capable of being expressed as all or a part of an antigenic polypeptide which is foreign to the host vector. In this embodiment, administration of the vaccine to an individual will result in augmentation and/or selective induction of the immune response of the individual to both antigenic polypeptide of the host vector and to co-expressed foreign antigenic polypeptide.

In another aspect, the present invention provides a method for producing an immune response in

a human or animal, in particular an immunodeficient or immunosuppressed human or animal, which comprises the step of administering to the human or animal a recombinant vaccine as broadly described above.

5 It will be appreciated from the broad description set out above that the present invention has particular application in the augmentation of immune responses in immunodeficient or immunosuppressed individuals. In one particularly
10 important aspect of this invention, there is provided a recombinant vaccine for use in the treatment of immunodeficient or immunosuppressed individuals infected with the human immunodeficiency virus (HIV), which comprises a vaccine vector which incorporates a
15 nucleotide sequence capable of being expressed as all or a part of an antigenic polypeptide derived from the human immunodeficiency virus (HIV), together with a second nucleotide sequence capable of being expressed as all or a portion of a lymphokine
20 effective in enhancing the immune response of the individual to the HIV antigenic polypeptide.

The lymphokines which may be expressed in vaccines according to this invention include those designated interleukin-1 (IL-1), interleukin-2
25 (IL-2), interleukin-3 (IL-3), interleukin-4 (IL-4) and γ -interferon (γ -IFN).

Interleukin-1 is a peptide hormone largely produced by activated macrophages. IL-1 modulates the proliferation, maturation and functional
30 activation of a broad spectrum of cell types (1-3) and plays a major role in the initiation and amplification of immune and inflammatory responses through its action on these diverse cell populations (4). The gene for murine (5) and human (6) IL-1 has

been cloned and expressed in E.coli.

Interleukin-2 is a lymphokine produced by helper T cells and is active in controlling the magnitude and type of the immune response (7). Other
5 functions have also been ascribed to IL-2 including the activation of NK cells (8) and the stimulation of cell division in large granular lymphocytes and B cells (9). Numerous studies in mice and humans have demonstrated that deficient immune responsiveness
10 both in vivo and in vitro can be augmented by IL-2. For example, exogenous IL-2 can restore the immune response in cyclophosphamide-induced immunosuppressed mice (10) and athymic (nude) mice (11). Furthermore, IL-2 can restore responsiveness of lymphocytes from
15 patients with various immunodeficiency states such as leprosy and cancer (12). IL-2 has also been used for the treatment of cancer (13). The gene for immune (14) and human (15) IL-2 has been cloned and sequenced.

20 Interleukin-3 is a hormone-like glycoprotein produced by lectin or antigen activated T lymphocytes and possibly other cells within the bone marrow. The hormone stimulates the growth and differentiation of haematopoietic progenitor cells and multipotential
25 stem cells and has been described under a variety of names, among them multi-colony stimulating factor, and haematopoietic growth factor (16). The gene for mouse IL-3 has been cloned and sequenced (17).

Interleukin-4 is a T cell derived factor
30 that acts as an induction factor on resting B cells, as a B cell differentiation factor and as a B cell growth factor (18). The factor also stimulates T cells and acts as a mast cell growth factor (18). The gene for murine (19) and human (20) IL-4 has been

isolated and sequenced.

γ -interferon is also a T cell derived molecule which has profound effects on the immune response. The molecule promotes the production of immunoglobulin by activated B cells stimulated with interleukin-2. γ -interferon also increases the expression of histocompatibility antigens on cells which associate with viral antigens to stimulate cytotoxic T cells. The gene for human γ -interferon has been isolated and sequenced (21).

The co-expression of a lymphokine such as IL-2 and an antigenic polypeptide by a recombinant vaccine (such as a recombinant virus vaccine) ensures that they are produced together by the same infected cells in a very localized area. This can be expected to lead to an elevation and acceleration of response to the virus vector component of the vaccine, e.g. vaccinia virus, with attendant benefits such as a reduction in the risks of complication associated with the use of vaccinia virus in normal individuals, and to those unidentifiable individuals who react adversely to vaccinia virus. Also, where there are immunological defects, as in the case of patients suffering from AIDS, leprosy or cytomegalovirus infection, co-expression of lymphokine could be instrumental in overcoming the defects to allow a normal response to the antigenic polypeptide and/or vector virus.

Furthermore, it is anticipated that the present invention will prevent or at least minimise the complications such as generalised vaccinia that can occur when vaccine is administered inadvertently to immunodeficient recipients (32):

[Further, cancer patients often show a

negligible or poor immunological response to the cancer antigens. It may be possible to enhance those responses to useful levels by taking cancer cells from the hosts, infecting them with, say, vaccinia virus/IL-2 recombinants, and returning them to the patient. To guard against generalized vaccinia infection or spread of the cancer cells it may, of course, be advisable to inactivate the recombinant-infected cancer cells prior to return to the patient.

Other lymphokines (e.g. IL-1, IL-3, IL-4 and γ -IFN), are involved in the control and augmentation of responses in other parts of the immune system including granulocyte-macrophage lineage, eosinophil differentiation and mucosal immunity. Construction of co-expressive vaccines will enable advantage to be taken of these specific modes of activity. Thus, as they are believed to have a role in the generation of protective responses at mucosal surfaces, such as in the gut, which promote expulsion of and immunity to parasites, a vaccine co-expressing IL-3 (or other lymphokine) with a helminth or other parasite antigenic polypeptide would be expected to give rise to enhanced immunity compared to that from the parasite antigen alone. Whilst a specific example of co-expression of the influenza haemagglutinin (HA) is described in detail herein, it will be appreciated that the present invention may be applied for the co-expression of other antigens including hepatitis virus, herpes simplex virus, Epstein-Barr virus and human immunodeficiency virus antigens, as well as malaria antigens.

As previously described, vaccinia virus has

been used as a vaccine vector to deliver antigens of unrelated infectious agents such as hepatitis B virus (22) and human immunodeficiency virus (23). The expression of an inserted gene in vaccinia virus
5 requires that the gene be placed next to a vaccinia promoter. The promoter usually used is designated p7.5 (24). This chimeric gene is then placed next to a DNA fragment of vaccinia virus taken from a non-essential region of the virus. Insertion into
10 infectious virus is by homologous recombination in which a marker rescue is used to select for virus recombinants. By way of example, the marker rescue can be either selection for thymidine kinase negative (TK^-) virus in which the foreign gene has been
15 inserted and thereby inactivating the TK gene; or by selecting for TK^+ virus in which the foreign gene is flanked by the herpes simplex TK gene. The latter is generally used to construct double recombinants that is, viruses expressing two foreign genes.

20 The expression of lymphokine genes in vaccinia virus may be detailed as two stages; the first is to create a plasmid in which the lymphokine is under the control of a vaccinia promoter 7.5 and downstream from a thymidine kinase (HSV) gene. This
25 plasmid is then used to transfect TK^- cells previously infected with a TK^- vaccinia virus expressing another foreign gene. TK^+ recombinant virus is then selected by culturing cells in the presence of methotrexate.

30 Although this invention has primarily been described with reference to vaccinia virus as the vaccine vector, it is to be understood that the inventive concept resides in co-expression of an antigenic polypeptide and lymphokine, and this

concept may be realized using other vaccine vectors, such as other poxvirus, herpes virus, adenovirus or bacteria.

It is also to be understood that the
5 invention is not limited by application to man or other species specifically mentioned herein, but may find application in a wide range of animal species.

Methods for construction and testing of recombinant vaccines according to this invention will
10 be well known to those skilled in the art, however, for better understanding of the invention some typical techniques will now be described. Standard procedures for endonuclease digestion, ligation and electrophoresis were carried out in accordance with
15 the manufacturer's or supplier's instructions. Standard techniques are not described in detail and will be well understood by persons skilled in the art.

EXAMPLE 1

Plasmids containing IL-1, IL-2, IL-3, IL-4
20 and γ -interferon are shown in Figures 1-5. Plasmids pmIL-11.270, pCD-JL-3, pCD-HuIL-2, pCD-IL-4 and pCD-Hu γ INF were obtained from DNAX Research Institute. The excised coding sequence is shown as the hatched bar. It is necessary to use different
25 restriction endonucleases to create suitable termini for insertion into plasmid; these are detailed in the respective diagrams. pBCB07 (25) contains the vaccinia 7.5K promoter, pFB-X (27) contains the HSV TK coding sequence (stippled) inserted at a BamH1
30 site downstream from a promoter in the vaccinia HindIII region. The recombinant plasmids pFB-X/IL1, pFB-X/IL2, pFB-X/IL3, pFB-X/IL4, pRB-X/ γ -interferon contain HSV-TK and lymphokine genes 3' to different vaccinia virus promoters and with flanking sequences.

derived from the HindIII F fragment of vaccinia virus. The orientations of genes and promoters are shown with arrows, vaccinia virus sequences with solid lines and plasmid DNA by thin lines. The pFB-
5 plasmids with inserted lymphokine genes are used to transfect 143B (TK⁻) cells previously infected with a TK⁻ vaccinia virus according to conditions previously described (24). The site of insertion of
10 HSV TK and lymphokine coding sequences and transposed vaccinia promoters in the vaccinia virus genome are shown in Figure 6. Vaccinia-virus WR strain HindIII restriction fragments are shown in the top line. Lower lines show in expanded form the DNA
15 configurations at insertion sites in the HindIII J and F fragments. Orientations of coding sequences and promoter sequences are shown with arrows.

EXAMPLE 2

This example describes the construction of a recombinant vaccinia virus (VV) expressing murine IL2
20 and the effect of the lymphokine on virus growth and immunogenicity.

Figure 6 shows:

(a) Genomic configuration of VV recombinants. A HindIII map of VV WR strain is shown
25 with insertion points at EcoRI (E) and BamHI (B) sites in the J and F fragments respectively. Arrows indicate orientations of VV TK gene, VV promoters and inserted influenza HA, HSV TK and murine IL2 coding sequences.

30 (b) Time course of IL2 production by VV-HA IL2-infected human 143B cells. IL2 activity in VV-HA-IL2-infected supernatants, circles and solid line; p VV-HA or uninfected cell supernatants, crosses and dotted line.

Figure 7 shows the growth of vaccinia virus recombinants in the foot pads of athymic Swiss outbred nude mice (a) and euthymic CBA/H mice (b). 2×10^7 PFU of VV-HA (triangles), VV-HA-TK (open circles) or VV-HA-IL2 (closed circles) were injected subcutaneously in 20 μ l into hind foot pads which were assayed for infectious virus on 143B cells on the indicated days. Points represent the titres of infectious virus present in individual mice.

As shown schematically in Figure 6a, cDNA encoding murine IL2 (14) was inserted into the HindIII F region of a VV recombinant, VV-HA (26), which already expressed the influenza haemagglutinin (HA). The IL2 recombinant virus, VV-HA-IL2, coexpressed HA and IL2 using the same VV 7.5Kd promoter but from separate sites in the viral genome. Since the herpes simplex virus (HSV) thymidine kinase (TK) gene was used as a selectable marker for virus construction a control virus VV-HA-TK, expressing HSV TK but not IL2 was constructed. Significant levels of biologically active IL2 were detected in supernatants from human 143B cells infected with VV-HA-IL2 within 4 hours and reached maximum activity around 12 hours (Figure 6b).

Athymic nude mice were inoculated into the right hind footpad with VV-HA-IL2 or control virus (VV-HA-TK). VV-HA-IL2 induced a mild swelling in the foot which resolved after several days; in contrast VV-HA-TK produced a severe necrotic lesion that remained unresolved for 30 days. After this time, high titres of virus (6×10^5 - 1.5×10^7 PFU) were recovered from the feet of the VV-HA-TK inoculated mice but not from mice given VV-HA-IL2. This suggested that the IL2 produced by the recombinant

virus enabled immunodeficient mice to control the virus infection. The kinetics of viral clearance from the feet of CBA/H mice were not significantly different for VV-HA-TK and VV-HA-IL2 (Figure 7a).

5 However, in nude mice, although titres of both VV-HA-TK and VV-HA-IL2 were high at day 3, indicating comparable rates of replication, VV-HA-IL2 was cleared by day 15 when no virus was detected in the feet. Titres of VV-HA-TK still remained high at day
10 15 (Figure 7b). Furthermore, when nude mice were injected intravenously (i.v.) with 10^6 PFU of VV-HA-IL2 or VV-HA-TK, mice given VV-HA-IL2 appeared unaffected by the virus whereas all mice given
15 this result infectious virus was recovered from spleens and lungs of VV-HA-TK- but not VV-HA-IL2-infected mice (Table 1).

TABLE 1. Vaccinia Virus recovered from Lungs
20 and Spleen

Mouse Strain	Organ	Log.10	Log.10
		<u>VV-HA-TK</u>	<u>VV-HA-IL2</u>
25 CBA/H	Lung	<2.0	<2.0
	Spleen	<2.0	<2.0
Outbred	Lung	4.91±0.27	<2.0
Nude	Spleen	3.64±0.11	<2.0

30

Lungs and spleens were collected from 3-4 mice/group 11 days post i.v. inoculation with 10^6 PFU of the indicated virus.

Methods

(a) Murine IL-2 cDNA was subcloned from pcD-IL-2, (14) generously provided by Dr K.I. Arai, DNAX Research Institute, Palo Alto, California. VV-HA-IL1 and VV-HA-TK were constructed by insertion
 5 of the HSV TK gene plus a chimeric promoter-IL2 fragment or alone, into the HindIII F region of the recombinant VV-HA (previously described as VV-PR8-HA6) (26) using plasmids as described in Example 1 and appropriate selection protocols (27,
 10 28).

(b) 143B cells 2×10^6 were infected with VV-HA-IL2 at 5 PFU/cell. The supernatants (1.5ml), harvested at the time points indicated, were assayed for IL2 activity using CTLL-2 cells (29) and the
 15 colourimetric method for cell growth of Mosmann (30). The results are presented as the log dilution of supernate producing 50% of maximum proliferation in the cell cultures.

EXAMPLE 3

20 A vaccinia virus expressing IL-3 was constructed as shown in Figure 2, using the methods described in Example 1.

Irradiated mice (650 Rads) injected with vaccinia virus (10^6 PFU intravenously) expressing
 25 IL-3 (VV-IL-3) show a reconstituted haemotopoietic system within seven days. Spleen cell counts are given below:

<u>Spleen Cell Number</u>			
		VV-IL-3	<u>NIL</u>
30	4 days	3×10^7	5×10^6
	7 days	2×10^8	2×10^6
	10 days	1×10^8	3×10^6

In addition, the injection of vaccinia virus expressing IL-3 (VV-IL-3) protects mice against the lethal effects of irradiation, as follows:

		<u>Deaths</u>	
5	<u>Irradiation Dose</u>	<u>VV-IL-3</u>	<u>NIL</u>
	950 Rads	0/6	6/6

EXAMPLE 4

10 This example describes the construction of a recombinant adenovirus expressing interleukin genes.

The starting virus for the adenovirus construct is adenovirus type 5 deletion mutant dl 327 that lacks the Xba fragment from 78.5 map units to 84.7 map units in early region 3 (31). This deletion
15 mutant allows the insertion of DNA without exceeding the amount of DNA that can be included in the virus particle. The removal of the E3 region also prevents production of a virus protein that complexes with the major histocompatibility heavy chain protein and
20 reduces the cell-mediated immune response to the virus. The Bam fragment from 60 map units to the right hand end of the viral DNA is cloned in plasmid. The plasmid DNA is cut downstream of the E3 promoter with a suitable restriction enzyme and the
25 interleukin gene inserted in place of the original E3 gene, under the control of the natural E3 promoter. The resulting plasmid containing the interleukin gene in the 60 to 100 map unit fragment of dl 327 is cut
30 with the appropriate restriction enzyme to separate viral and plasmid DNA and transfected into cells together with the overlapping EcoR1 A fragment (0 to 76 map units) of wild type virus. Recombination between the two overlapping DNA fragments will reconstitute viable adenovirus in which the E3 gene

is replaced by the interleukin gene.

EXAMPLE 5

Figure 8 outlines the construction of human immunodeficiency virus IL-2 recombinant vaccinia virus in accordance with the present invention. 5 pFB-X/IL-2 is constructed as shown using the methods described in Example 1. The construction of pTG1125 is as previously described (35).

REFERENCES

1. Howard, M. et al. J. Exp. Med. 157, 1529-1534 (1983).
2. Baracos, V. et al. New Eng. J. Med. 308, 553-558 (1983).
3. Schmidt, J.A. et al. J. Immunol. 128, 2177-2182 (1982).
4. Mizel, S.B. in Biological Response in Cancer (ed. Mihich E.) 89-119 (Plenum, New York 1982).
5. Lomedico, P.T. et al. Nature 312, 458-461 (1984).
6. Auron, P.E. et al. Proc. Natl. Acad. Sci. US 81, 7907-7911 (1984).
7. Smith, K.A. Ann. Rev. Immunol. 2, 319-333 (1984).
8. Minato, N. et al. J. Exp. Med. 154, 750 (1983).
9. Tsudo, M. et al. J. Exp. Med. 160, 612-616 (1984).
10. Merluzzi, V.J. et al. Cancer Res. 41, 850-853 (1981).
11. Wagner, H. et al. Nature 284, 278-80 (1982).
12. Vose, B.M. et al. Cancer Immuno. 13, 105-111 (1984).
13. Rosenberg, S.A. et al. Science 233, 1318-1321 (1986).
14. Yokota, T. et al. Proc. Natl. Acad. Sci. USA 82, 68-72 (1985).
15. Taniguchi, T. et al. Nature, 302, 305-307 (1983).
16. Schroder, T.W. et al. in Contemporary Topics in Molecular Immunology: The interleukins, eds Gillis, S. and Inman, F.P. (Plenum, N.Y.) Vol. 10, pp. 121-146 (1985).
17. Yokota, T. et al. Proc. Natl. Acad. Sci. USA 81, 1070-1074 (1984).
18. Sevenusar, E. Eur. J. Immunol. 17, 67-72 (1987).
19. Lee, F. et al. Proc. Natl. Acad. Sci. USA 83,

- 2061-2065 (1986).
20. Yokota, T. et al. Proc. Natl. Acad. Sci. USA 83, 5894-5898 (1986).
21. Gray, P.W. et al. Nature 295, 503-508 (1982).
22. Smith, G.L. et al. Nature 302, 490-495 (1983).
23. Hu, S.L. et al. Nature 320, 537-540 (1986).
24. Boyle, D.B. and Coupar, B.E.H. J. Gen. Virol. 67, 1591-1600 (1986).
25. Boyle, D.B. et al. Gene 35, 169-177 (1985).
26. Andrew, M.E. et al. Microbial Pathogenesis 1, 443-452 (1986).
27. Coupar, B.E.H. et al. J. Gen. Virol. 68, (1987), (In press).
28. Coupar, B.E.H. et al. Gene (In preparation).
29. Gillis S. et al. J. Immunol 120, 2027-2032 (1978).
30. Mosmann, J., J. Immunol. Meth. 65, 55-63 (1983).
31. Cutt, J.R. et al. J. Virol. 61, 543 (1987).
32. Wright, D.C. et al, New Eng. J. Med. 316, 673-675 (1987).
33. Kierny, M.P. et al. Biotechnology 4, 790-795 (1986).

CLAIMS:

1. A recombinant vaccine comprising a vaccine vector which incorporates a first nucleotide sequence capable of being expressed as all or a part of an antigenic polypeptide, together with a second nucleotide sequence capable of being expressed as all or a part of a lymphokine effective in enhancing the immune response to the antigenic polypeptide.
2. A recombinant vaccine according to claim 1, wherein said vaccine vector incorporates a nucleotide sequence capable of being expressed as all or a part of an antigenic polypeptide which is foreign to the host vector.
3. A recombinant vaccine according to claim 1 or claim 2, wherein said vaccine vector is a virus.
4. A recombinant vaccine according to claim 3, wherein said virus is a poxvirus, herpes virus or adenovirus.
5. A recombinant vaccine according to claim 4, wherein said virus is vaccinia virus.
6. A recombinant vaccine according to any one of claims 1 to 5, wherein said lymphokine is selected from the group consisting of interleukin-1, interleukin-2, interleukin-3, interleukin-4 and γ -interferon.
7. A method for the production of a recombinant vaccine according to any one of claims 1 to 6, which

comprises the step of inserting into a vaccine vector a nucleotide sequence capable of being expressed as all or a part of a lymphokine.

8. A method according to claim 7, which comprises the further step of inserting into the vaccine vector a nucleotide sequence capable of being expressed as all or a part of an antigenic polypeptide which is foreign to the host vector.


9. Use of a recombinant vaccine according to any one of claims 1 to 6, for the production of an immune response in a human or animal, particularly an immunodeficient or immunosuppressed human or animal.


10. A method for producing an immune response in a human or animal, particularly an immunodeficient or immunosuppressed human or animal, which comprises the step of administering to the human or animal a recombinant vaccine according to any one of claims 1 to 6.

11. A recombinant vaccine for use in the treatment of immunodeficient or immunosuppressed individuals infected with the human immunodeficiency virus (HIV), which comprises a vaccine vector which incorporates a nucleotide sequence capable of being expressed as all or a part of an antigenic polypeptide derived from the human immunodeficiency virus (HIV), together with a second nucleotide sequence capable of being expressed as all or a part of a lymphokine effective in enhancing the immune response of the individual to the HIV antigenic polypeptide.


12. A recombinant vaccine according to claim 11, wherein said vaccine vector is vaccinia virus.


13. A recombinant vaccine according to claim 11 or claim 12, wherein said lymphokine is selected from the group consisting of interleukin-1, interleukin-2, interleukin-3, interleukin-4 and γ -interferon.


**FIG 1**




2/9

**FIG 2****SUBSTITUTE SHEET**



FIG 3

SUBSTITUTE SHEET






FIGURE 4




FIGURE 5




FIG 6a







FIG 6b






8/9

FIG 7









FIG 8

INTERNATIONAL SEARCH REPORT

International Application No. PCT/AU 87/00246

I. CLASSIFICATION OF SUBJECT MATTER : 1. 50-0-0-0 Classification Symbols apply, indicate only 1		
According to International Patent Classification (IPC) or to both National Classification and IPC		
Int. Cl. ⁴ C12N 15/00, 7/00, A61K 39/285, 39/21, 39/145, 45/02, 37/02 / C12R 1:91		
II. FIELDS SEARCHED		
Minimum Documentation Searched *		
Classification System	Classification Symbols	
IPC	C12N 15/00	
Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in the Fields Searched *		
AU : IPC as above C.A. KEYWORDS (INTERFERON or INTERLEUKIN) and (VACCINIA or ADENOVIRUS)		
III. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of Document ** with indication, where appropriate, of the relevant passages **	Relevant to Claim No. **
X	AU,A, 36635/84 (GENENTECH INC.) 20 June 1985 (20.06.85) See page 11	(1-3,6-8)
X,P	EP,A, 206939 (TRANSGENE S.A.) 30 December 1986 (30.12.86) (& AU 59676/86)	(1,3-7)
X,P	EP,A, 206920 (TRANSGENE S.A.) 30 December 1986 (30.12.86) (& AU 59579/86)	(1,3-7)
X,P	AU,A, 59453/86 (KANEGAFUCHI KAGAKU KOGYO KABUSHIKI KAISHA) 8 January 1987 (08.01.87) See page 23 lines 11-19	(1,3,7)
X	AU,A, 11674/83 (BIOGEN N.V.) 1 September 1983 (01.09.83) See page 25 lines 1-23	(1,3,6,7)
X	Proceedings of the National Academy of Science USA, Volume 82, No.3 issued 1985, February (Washington D.C.), R.J. Kaufman, "Identification of the components necessary for adenovirus translational control and their utilization in cDNA expression vectors", see pages 689-693	(1,3,4,6,7)
<div style="display: flex; justify-content: space-between;"> <div style="width: 48%;"> <p>* Special categories of cited documents: **</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="width: 48%;"> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"G" document member of the same patent family</p> </div> </div>		
IV. CERTIFICATION		
Date of the Actual Completion of the International Search		Date of Mailing of this International Search Report
17 November 1987 (17.11.87)		(30-11-87) 30 NOVEMBER 1987
International Searching Authority Australian Patent Office		Signature of Authorized Officer  J.H. CHAN

ANNEX TO THE INTERNATIONAL SEARCH REPORT ON
INTERNATIONAL APPLICATION NO. PCT/AU 87/00246

This Annex lists the known "A" publication level patent family members relating to the patent documents cited in the above-mentioned international search report. The Australian Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

Patent Document Cited in Search Report		Patent Family Members			
AU 36635/84		DK 6009/84	EP 146354	FI 844934	
		HU 36187	IL 73803	JP 60202899	
		NO 845042	PT 79691	ZA 8409767	
EP 206920		AU 59579/86	DK 798/87	FR 2583429	
		WO 8607609	ZA 8604525		
EP 206939		AU 59676/86	FR 2583770	ZA 8604561	
		DK 878/87			
AU 59453/86		EP 207518	JP 62011095		
AU 11674/83		DK 734/83	EP 88540	ES 519953	
		IL 67961	JP 59051792	ZA 8301094	

END OF ANNEX

0. 88/00971(1) ()
12N15/85-(C12N15/36)-(C12N15/66)-
A61K39/12)-

0 275 300

MENTARY

AN SEARCH REPORT

Application number

--- BASIC DOC.---

C12N15/85

the European Patent Convention
the purposes of subsequent

EP 87 90 5142

proceedings, as the European search report

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.4)
X	EP-A-0 181 117 (AMERICAN HOME PRODUCTS CORP.) * Whole document * --	4	C 12 N 15/00 C 12 N 7/00 A 61 K 39/285 A 61 K 39/21 A 61 K 39/145 A 61 K 45/02 A 61 K 37/02//
X	EP-A-0 176 170 (INSTITUT MERIEUX) * Whole document * --	4	C 12 R 1:91
X	ADV. IMMUNOPHARMACOL. vol. 3, Proc. Int. Conf. 3rd, 1985, (publ. 1986), pages 223-230, Ed. Pergamon Press, Oxford, GB C.A. DINARELLO: "The biological activities of human interleukin-1: purified and recombinant materials" * Page 228, lines 1-13 * --	1,2 6-8, 11-13	TECHNICAL FIELDS SEARCHED (Int. Cl.4)
Y			
Y	BIOLOGICAL ABSTRACTS, RRM no. 29067133 H. KAWAMURA et al.: "Immunization with antigen and interleukin 2 . / .		C 12 N
INCOMPLETE SEARCH			
<p>The Search Division considers that the present European patent application does not comply with the provisions of the European Patent Convention to such an extent that it is not possible to carry out a meaningful search into the state of the art on the basis of some of the claims.</p> <p>Claims searched completely: 1-8, 11-13 Claims searched incompletely: 9, 10 Claims not searched: Reason for the limitation of the search:</p> <p>Method for treatment of the human or animal body by surgery or therapy (See art. 52(4) of the European Patent Convention)</p>			
Place of search The Hague		Date of completion of the search 16-10-1988	Examiner SKELLY
CATEGORY OF CITED DOCUMENTS			
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000



DOCUMENTS CONSIDERED TO BE RELEVANT			CLASSIFICATION OF THE APPLICATION (Int. Cl.4)
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	
	in-vivo overcomes IR gene low responsiveness" & JOURNAL OF EXPERIMENTAL MEDICINE (US), 1985, vol.162, no. 1, p. 381-386 * Title and terms * --	6-8,13	
Y	CHEMICAL ABSTRACTS, vol. 92, no. 19, May 12, 1980, page 446, abstract 162033q, Columbus, Ohio, US T.A. BEKTEMIROV et al.: "Smallpox vaccination under the protection of interferon and its inducer" & VOPR.VIRUSOL. 1980, (1), 76-8 * Abstract * --	6-8,13	TECHNICAL FIELDS SEARCHED (Int. Cl.4)
Y	EP-A-0 138 133 (SCHERING BIOTECH. CORP.) * Whole document * --	6-8,13	
Y	PROC. NATL. ACAD. SCI. USA; vol.83, no. 16, August, 1986, pages 5894-5898, Washington, US T. YOKOTA et al.: "Isolation and characterization of a human interleukin cDNA clone, homologous to mouse B-cell stimulatory factor 1, that expresses B-cell- and T-cell-stimulating activities" * Whole document * --	6-8,13	
Y	NATURE; vol. 320, no. 6062, April 1986, London, GB; pages 537-540 S.-L. HU et al.: "Expression of AIDS virus envelope gene in recombinant vaccinia viruses" * Whole document * --	11-13	
X	THE YEARBOOK OF AGRICULTURE, US Department of Agriculture, 1986, pages 104-108 K.J. CREMER: "New vaccines for old diseases" * Page 107 *	1-8	

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100



DOCUMENTS CONSIDERED TO BE RELEVANT			CLASSIFICATION OF THE APPLICATION (Int. Cl. 4)
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	
Y	--	11-13	
A	BIOLOGICAL ABSTRACTS, vol. 79, 1985, no. 13273 P.J. GROB et al: "Interferon as an adjuvant for hepatitis B vaccination in non-responder and low-responder populations" & EUR. J. CLIN. MICROBIOL. 3(3), 195-109 * Abstract *	6-8,13	
A	-- J. IMMUNOL., vol. 130, no. 5, May 1983, pages 2191-2194 M.J. STARUCH et al.: "The adjuvanticity of interleukin 1 in vivo" * Whole document *	6-8,13	TECHNICAL FIELDS SEARCHED (Int. Cl. 4)
T	-- NATURE, vol. 329, no. 6139, 8-14th October 1987, London, GB pages 545-546 I.A. RAMSHAW et al.: "Recovery of immunodeficient mice from a vaccinia virus/IL-2 recombinant infection" * Whole document *	1	

100

1

100

1

1

1

1

1

1

1

1

1

1

CLAIMS:

1. A recombinant vaccine comprising a vaccine vector which incorporates a first nucleotide sequence capable of being expressed as all or a part of an antigenic polypeptide, together with a second nucleotide sequence capable of being expressed as all or a part of a lymphokine effective in enhancing the immune response to the antigenic polypeptide.
2. A recombinant vaccine according to claim 1, wherein said vaccine vector incorporates a nucleotide sequence capable of being expressed as all or a part of an antigenic polypeptide which is foreign to the host vector.
3. A recombinant vaccine according to claim 1 or claim 2, wherein said vaccine vector is a virus.
4. A recombinant vaccine according to claim 3, wherein said virus is a poxvirus, herpes virus or adenovirus.
5. A recombinant vaccine according to claim 4, wherein said virus is vaccinia virus.
6. A recombinant vaccine according to any one of claims 1 to 5, wherein said lymphokine is selected from the group consisting of interleukin-1, interleukin-2, interleukin-3, interleukin-4 and γ -interferon.
7. A method for the production of a recombinant vaccine according to any one of claims 1 to 6, which

comprises the step of inserting into a vaccine vector a nucleotide sequence capable of being expressed as all or a part of a lymphokine.

8. A method according to claim 7, which comprises the further step of inserting into the vaccine vector a nucleotide sequence capable of being expressed as all or a part of an antigenic polypeptide which is foreign to the host vector.

9. Use of a recombinant vaccine according to any one of claims 1 to 6, for the production of an immune response in a human or animal, particularly an immunodeficient or immunosuppressed human or animal.

10. A method for producing an immune response in a human or animal, particularly an immunodeficient or immunosuppressed human or animal, which comprises the step of administering to the human or animal a recombinant vaccine according to any one of claims 1 to 6.

11. A recombinant vaccine for use in the treatment of immunodeficient or immunosuppressed individuals infected with the human immunodeficiency virus (HIV), which comprises a vaccine vector which incorporates a nucleotide sequence capable of being expressed as all or a part of an antigenic polypeptide derived from the human immunodeficiency virus (HIV), together with a second nucleotide sequence capable of being expressed as all or a part of a lymphokine effective in enhancing the immune response of the individual to the HIV antigenic polypeptide.

12. A recombinant vaccine according to claim 11, wherein said vaccine vector is vaccinia virus.

13. A recombinant vaccine according to claim 11 or claim 12, wherein said lymphokine is selected from the group consisting of interleukin-1, interleukin-2, interleukin-3, interleukin-4 and γ -interferon.

